

Jurnal Ilmiah
FARMASI

(Scientific Journal of Pharmacy)



JURNAL ILMIAH FARMASI
(SCIENTIFIC JOURNAL OF PHARMACY)

PIMPINAN UMUM/ PENANGGUNG JAWAB
Dekan Fakultas Matematika dan Ilmu Pengetahuan Alam
Universitas Islam Indonesia

WAKIL PIMPINAN UMUM/ WAKIL PENANGGUNG JAWAB
Ketua Jurusan Farmasi FMIPA UII

MITRA BESTARI

1. Prof. Dr. Wiryatun Lestariana, Apt
2. Prof. Dr. Zullies Ikawati, Apt
3. Prof. Dr. Sudibyo Martono, Apt
4. Dr. Tedjo Yuwono, Apt
5. Prof. Dr. Dachriyanus, Apt
6. Prof. dr. Iwan Dwiprahasto, MMedSc, PhD
7. Prof. Dr. Lukman Hakim M.Sc., Apt
8. Prof. Dr. Achmad Fudholi, DEA, Apt
9. Prof. Dr. Ibnu Gholib Gandjar, DEA., Apt

DEWAN EDITOR

Ketua : Saepudin, M.Si., Apt
Sekretaris : Rochmy Istikharah, M.Sc., Apt.
Anggota : Vitarani Dwi Ananda Ningrum, M.Si., Apt
Okti R. Mafruhah, MSc., Apt
Dimas Adhi Pradana, MSc., Apt.
Fithria DA. Suryanegara, MSc., Apt.
Ari Wibowo, S.Farm., Apt
Arba Pramudita Ramadani, MSc., Apt.
Oktavia Indrati, S.Farm., Apt.

Penerbit

Jurusan Farmasi Fakultas Matematika dan Ilmu Pengetahuan Alam
Universitas Islam Indonesia

Alamat Penerbit

Jurusan Farmasi FMIPA UII
Jl. Kaliurang Km. 14,4 Yogyakarta 55584
Telp. (0274) 896439 ext. 3047
Email: jif@uii.ac.id

BIOACTIVE CONSTITUENTS FROM THE LEAVES OF *MELASTOMA MALABATHRICUM* L.

Deny Susanti^{1*}, Hasnah M. Sirat², Farediah Ahmad³, Rasadah Mat Ali³

¹Department of Biomedical Science, Faculty of Science, International Islamic University Malaysia, 25200 Kuantan, Malaysia.

²Department of Chemistry, Faculty of Science, Universiti Teknologi Malaysia, 81310 Skudai, Johor, Malaysia

³Medicinal Plant Division, Forest Research Institute Malaysia, 52109 Kepong, Kuala Lumpur, Malaysia

*e-mail: deny@iiu.edu.my

ABSTRACT

Phytochemical and bioactivity studies of the leaves of *Melastoma malabathricum* L. (Melastomataceae) have been investigated. The *n*-hexane extract yielded α -amyrin, patriscabatrane and auranamide, ethyl acetate extract gave quercetin and quercitrin, and methanol extract gave quercitrin and kaempferol-3-*O*-(2'',6''-di-*O-p-trans*-coumaroyl)glucoside. The crude extracts and isolated compounds were screened for their antioxidant and cytotoxic activities. The antioxidant assay was carried out by FTC and DPPH radical scavenging method. Kaempferol-3-*O*-(2'',6''-di-*O-p-trans*-coumaroyl)glucoside, quercetin and quercitrin showed strong activities with inhibition more than 90% in the FTC method. Quercetin was found to be the most active as radical scavenger in DPPH method with IC₅₀ of 0.69 μ M. α -Amyrin and kaempferol-3-*O*-(2'',6''-di-*O-p-trans*-coumaroyl)glucoside demonstrated the strongest activities in the anti-inflammatory assay of TPA mouse ear oedema with IC₅₀ of 0.11 and 0.34 mM/ear, respectively.

Keywords: Antiinflammatory, Antioxidant, Flavonoids, Melastomataceae, *Melastoma malabathricum*, Terpenoid